


# Bc at CDF




William Wester  
Fermilab  
for the CDF collaboration




# Introduction

## B Physics at Hadron Colliders

- UA1 cross section measurements
- CDF fully reconstructed  $B \rightarrow J/\psi K^{(*)}$



UA1  $\sigma(b)$  in  $\mu$  channel  
PLB 213, 415 (1988)



CDF  $B_u \rightarrow J/\psi K$   
PRL 68, 3403 (1992)

Since the 1980's ...

### Advantages:

- Large  $\sigma(b) \times \mathcal{L}$
- All mesons and baryons
- Triggerable:  $\ell$  or  $J/\psi$
- Multipurpose detectors





### Disadvantages: (perceived)

- High backgrounds
- Limited acceptance
- Small Lorentz boost
- Unknown initial state

- Study of  $B_c$  highlights hadron collider advantages
  - Large cross section for producing triggerable low background decays not accessible at the  $B$  factories.




# Tevatron and CDF in Run II





# Bc properties


- Bc is a heavy-heavy system
  - Production: Factorization with two scales  $M_b + M_c$  and contributions of color singlet / octet
    - Softer  $P_T$  distribution?
  - Decay: both b and c quarks can participate
    - Shorter *c-like* lifetime?
    - Large number of final state *BRs*.
  - Mass: new system for potential models and new lattice QCD calculations
- All aspects of the theoretical work require experimental measurement => happening now at the Tevatron





# Bc in Run I ('91-'96)

- A few candidate events at LEP and the CDF observation and measurements...




PRL 81, 2432 (1998) and PRD 58, 112004 (1998)

Production measurement ( $P_T(B) > 6 \text{ GeV}/c$   $|\eta| < 0.6$ ):

$$\frac{\sigma(B_c) \times B(B_c \rightarrow J/\psi \ell v)}{\sigma(B_u) \times B(B_u \rightarrow J/\psi K)} = 0.132^{+0.041}_{-0.037} \text{ (stat)} \pm 0.031 \text{ (syst)} \quad c\tau^{+0.032}_{-0.020} \text{ (} c\tau \text{)}$$

Note: assuming harder  $P_T$  spectrum in MC





## Run II results: semi-leptonic decays

---

- $B_c \rightarrow J/\psi + \ell\nu$  with  $\ell = e, \mu$
- Not fully reconstructed (missing  $\nu$ )
- Understanding backgrounds are key
  - $b\bar{b}$  events with the  $J/\psi$  from b and  $\ell$  from  $\bar{b}$
  - Fake muons or fake electrons
  - Other backgrounds
- Study  $J/\psi + \text{track}$  and  $B_u \rightarrow J/\psi K$
- Look for  $B_c$  excess above background and make measurements




# B<sub>c</sub> → J/ψ μ X

- Use 2.7M J/ψ's in 0.36 fb<sup>-1</sup>
- Combine with third track with & w/o muon ID
  - $P_T > 3 \text{ GeV}$ ,  $c\tau > 60\mu\text{m}$ , and  $\Delta\phi(\text{J}/\psi - \text{trk}) < 90 \text{ deg}$
- Use B<sub>u</sub>→J/ψK from data for normalization
- Use Monte Carlo of B<sub>u</sub> and B<sub>c</sub> for  $\epsilon_{\text{rel}}$
- Evaluate backgrounds in the data
  - Fake muons, b $\bar{b}$ , fake J/ψ
- Estimate systematic uncertainties
- Fit data in 4-6 GeV for signal and backgrounds
  - Evaluate relative production of B<sub>c</sub> to B<sub>u</sub>




# Fake muon background






How many come from  
 $J/\psi + \text{track}$  where the  
track is a fake muon?

Fake muons primarily from  
decay in flight:  $16.3 \pm 2.9$   
estimated in  $4 < M < 6 \text{ GeV}$ .






- Fake muons: determine  $\pi$ ,  $K$ ,  $p$  composition vs  $P_T$  ( $dE/dx$  and TOF) and then use  $D^*$ ,  $\Lambda$  decays to find fakes vs  $P_T$





# More backgrounds

- bb background from Pythia Monte Carlo normalized to  $B_u \rightarrow J/\psi K$  data using  $\Delta\phi$  distributions (vary production)
- Fake  $J/\psi$  from  $J/\psi$  sidebands




Backgrounds from the other b:  $12.7 \pm 1.7 \pm 5.7$  estimated in  $4 < M < 6$  GeV.

Backgrounds from fake  $J/\psi$  (no double counting):  $19.0 \pm 3.0$  estimated in  $4 < M < 6$  GeV.



# Muon channel results

Mass window	3.0 – 4.0 GeV/c <sup>2</sup>	4.0 – 6.0 GeV/c <sup>2</sup> (signal)	6.0 – 10.0 GeV/c <sup>2</sup>
$B_c$ candidates in mass window	$7 \pm 2.4$	$106 \pm 10.3$	$19 \pm 4.2$
Fake muon background	$3.9 \pm 0.7$	$16.3 \pm 2.9$	$2.2 \pm 0.4$
$BB$ background	$0.6 \pm 0.4 \pm 0.1$	$12.7 \pm 1.7 \pm 5.7$	$6.0 \pm 1.1 \pm 1.8$
Fake $J/\psi$ background	$0.5 \pm 0.5$	$19.0 \pm 3.0$	$5.0 \pm 1.7$
Fake $\mu$ from ( $J/\psi_{side} + Trk$ )	$0.3 \pm 0.1$	$2.0 \pm 0.5$	$0.7 \pm 0.2$
Total Background	$4.7 \pm 0.9$	$46.0 \pm 7.3$	$12.5 \pm 2.7$
Events above background	$2.5 \pm 2.8$	$60.0 \pm 12.6$	$6.5 \pm 5.1$



Use MC for relative efficiency for  $B_c$  and  $B_u$  along with  $B_u \rightarrow J/\psi K$  to obtain:

$$\frac{\sigma(B_c) \times \mathcal{B}(B_c \rightarrow J/\psi \ell v)}{\sigma(B_u) \times \mathcal{B}(B_u \rightarrow J/\psi K)} = \begin{array}{l} P_T(B) > 4 \text{ and} \\ |\eta| < 1 \end{array}$$

$$0.249 \pm 0.045 \pm 0.069 \pm 0.082$$


Other measurements from this sample are in preparation.

CDF Note: 7649



# Bc → J/ψ e X


- Fake electron
  - Use J/ψ+track data
  - Estimate fake rate from data ( $D^0 \rightarrow K\pi, \Lambda^0 \rightarrow p\pi$ )
- Photon conversion
  - Use J/ψ+tagged conversion data
  - Conversion finding efficiency from MC
- $b\bar{b}$  background
  - $b \rightarrow J/\psi X$  and  $\bar{b} \rightarrow e X$
  - PYTHIA  $b\bar{b}$  Monte Carlo





# Photon conversions

- Remove conversions by finding the partner track during the electron selection
- Evaluate the conversion finding efficiency from MC
- Calculate the residual conversion background as a function of  $M(J/\psi e)$  using  $J/\psi +$ tagged conversions.
- Expected background
  - $14.54 \pm 4.38(\text{stat}) \pm 6.39 (\text{syst})$






# electron channel results

- Observed  
 $178.5 \pm 14.7$  events
- Background  
 $63.6 \pm 4.9 \pm 13.6$
- Excess  
 $114.9 \pm 15.5 \pm 13.6$
- Significance  
 $5.9\sigma$

$$\frac{\sigma(B_c) \times B(B_c \rightarrow J/\psi \ell v)}{\sigma(B_u) \times B(B_u \rightarrow J/\psi K)} =$$

$P_T(B) > 4$  and  
 $|y| < 1$

$0.282 \pm 0.038(\text{stat.}) \pm 0.035(\text{yield}) \pm 0.065(\text{acceptance})$






# Precise lifetime measurement

- Release e sample  $c\tau$  cuts
- Evaluate prompt background
- Determine K-factors (missing- $\nu$ )
  - Difference between pseudo- $c\tau$  and  $c\tau$
- Fit for lifetime: signal and background
- Evaluate systematics


K-factors vs  $M(J/\psi e)$





# Bc lifetime

CDF Note: 7758 CDF Run 2 Preliminary :  $\sim 360 \text{ pb}^{-1}$



## Systematic uncertainties

catalog	description	Fitted $c\tau (\mu\text{m})$	$\Delta c\tau (\mu\text{m})$
K-factor	$M(B_c) = 6.4, 6.2 \text{ GeV}$	$140.4^{+21.7}_{-19.5}, 143.0^{+22.0}_{-19.8}$	$\pm 1.7$
K-factor	$\tau(B_c) = 0.5, 0.7 \text{ ps}$	$141.9^{+21.9}_{-19.6}$	$\pm 0.2$
K-factor	$H_b \rightarrow J/\psi X$ spectrum	$140.8^{+21.7}_{-19.5}$	$\pm 1.3$
K-factor	Inclusive $J/\psi X e\nu$	$140.5^{+21.7}_{-19.5}$	$\pm 1.6$
K-factor	trigger simulation	$142.4^{+21.9}_{-19.7}$	$\pm 0.3$
K-factor sub-total $\Delta c\tau = \pm 2.7$			
$\mathcal{F}_{fake-e}$	Use $J/\psi + \text{trk}$ shape directly	$140.6^{+21.5}_{-19.4}$	$-1.5$
$\mathcal{F}_{fake-J/\psi}$	use $J/\psi + e$ sideband	$136.0^{+21.8}_{-22.6}$	$-6.1$
$\mathcal{F}_{conv-e}$	Use tagged conv-e shape directly	$141.2^{+21.7}_{-19.5}$	$-0.9$
$\mathcal{F}_{conv-e}$	use $J/\psi + \text{conv-e}$ sideband	$144.8^{+21.5}_{-19.3}$	$+2.7$
$\mathcal{F}_{bb}$	use FE only	$150.2^{+17.5}_{-15.9}$	$+8.1$
$\mathcal{F}_{bb}$	use GS only	$138.3^{+16.6}_{-15.0}$	$-3.8$
$\mathcal{F}_{bb}$	No error scaling in MC	$140.9^{+21.6}_{-19.5}$	$-1.2$
Background shapes sub-total $\Delta c\tau = (+8.5, -7.5)$			
$L_{xy}$ resolution	extra Gaussian/symmetric exponential	$137.0^{+21.9}_{-19.8}, 136.5^{+21.9}_{-19.8}$	$-5.6$
$L_{xy}$ resolution	Punzi effect	$137.3^{+22.1}_{-19.4}$	$-4.8$
$L_{xy}$ resolution	silicon alignment		$\pm 1$
$L_{xy}$ resolution sub-total $\Delta c\tau = (+1.0, -7.4)$			
Total systematic error $\Delta c\tau = (+9.0, -10.9)$			


$$c\tau(B_c) = 142.1^{+21.9/-19.7(\text{stat}) \pm 10.0(\text{syst})} \mu\text{m}$$

$$\tau(B_c) = 0.474^{+0.073/-0.066(\text{stat}) \pm 0.033(\text{syst})} \text{ ps}$$






# Bc → J/ψ π

- Full reconstruction allows for precise mass measurement
- New analysis
  - Tune selection on the data:  
 $Bu \rightarrow J/\psi K$  reference decay
  - After approval, "open box".
  - Wait for events to become a significant excess
  - Measure properties of the  $B_c$





# Bc → J/ψ π





Num(events)<sub>FIT</sub> =  
38.9 sig 26.1 bkg  
between 6.24-6.3  
Significance > 6σ  
over search area

$$\text{Mass}(B_c) = 6275.2 \pm 4.3 \pm 2.3 \text{ MeV}/c^2$$



# Recent Lattice Calculations

- Lattice calculations that show good agreement with experiment were used to *predict* the mass of the  $B_c$



$$M(B_c)_{\text{CDF}} = 6275.2 \pm 4.3 \pm 2.3 \text{ MeV}/c^2$$

$$M(B_c)_{\text{LAT}} = 6304 \pm 12^{+18}_{-0} \text{ MeV}/c^2$$



# Summary and conclusions

---

- The study of the  $B_c$  is happening in Run II
- Semi-leptonic decays observed  $>5\sigma$ 
  - $J/\psi \mu X$  and  $J/\psi e X$
  - Precise lifetime measured
  - $\tau(B_c) = 0.474 +0.073/-0.066(\text{stat}) \pm 0.033(\text{syst}) \text{ ps}$
- Fully reconstructed  $B_c \rightarrow J/\psi \pi$  sample  $>6\sigma$ 
  - Precision mass compared with theory
  - $M(B_c) = 6275.2 +/- 4.3(\text{stat}) +/- 2.3(\text{syst}) \text{ MeV}/c^2$
- $B_c$  at CDF is challenging theory with promise of more to come