



Combination of CDF top quark mass measurements using up to 5.6 fb^{-1} of data

The CDF Collaboration
URL <http://www-cdf.fnal.gov>
(Dated: July 5, 2010)

We summarize the CDF measurements of the top-quark mass, M_{top} , performed analyzing 110 pb^{-1} of data collected at 1.8 TeV center of mass collision energy, and as much as 5.6 fb^{-1} of data collected at 1.96 TeV. We combine the most precise CDF published and preliminary results to obtain $M_{top} = 173.13 \pm 0.67 \text{ (stat)} \pm 0.95 \text{ (syst)} \text{ GeV}/c^2$ which corresponds to a total uncertainty of $1.16 \text{ GeV}/c^2$. The combination provides a relative precision $\Delta M_{top}/M_{top}$ of 0.67%. Rounding off to two significant digits, the combination yields $M_{top} = 173.1 \pm 1.2 \text{ (stat + syst)} \text{ GeV}/c^2$.

Preliminary Results for Summer 2010 Conferences

I. INTRODUCTION

We combine the CDF published top-quark mass results from the Tevatron run at 1.8 TeV center of mass collision energy (Run I) performed analyzing 110 pb^{-1} of data [1-3] with two published [4,5] and two preliminary results [6,7] that use up to 5.6 fb^{-1} of data collected at the Tevatron run at 1.96 TeV center of mass energy (Run II).

Results from the $t\bar{t} \rightarrow qq' bqq'\bar{b}$ (HAD), $t\bar{t} \rightarrow \ell\nu qq' b\bar{b}$ (LJT), and $t\bar{t} \rightarrow \ell^+\nu b\ell^-\bar{b}$ (DIL) final states are included. These measurements are combined accounting for statistical and systematic correlations using the method of reference [8].

Relative to the previous CDF combination reported in [9], this combination includes updates of the Run II analyses in the LJT and DIL channels. The recent Run II measurement in the LJT channel using Matrix Element technique yields the single most precise result, with a 0.72% relative precision on $\Delta M_{top}/M_{top}$, more precise than the '09 Tevatron top mass combination [10]. A new measurement in the DIL channel provides slightly better precision than the one used in the previous combination.

The uncertainty categories used in the combination are detailed in Section II while the input measurements themselves are summarized in Section III. The correlations used in the combination are discussed in Section IV and the resulting top-quark mass is given in Section V. Section VI summarizes the results obtained.

II. UNCERTAINTY CATEGORIES

We employ the same uncertainty categories as used for the Tevatron world average [10]. They have evolved to include a detailed breakdown of the various sources of uncertainty and aim to lump together sources of systematic uncertainty that share the same or similar origin. For example, the ‘‘Signal’’ category discussed below includes the uncertainties from initial state radiation (ISR), final state radiation (FSR), and parton distribution functions (PDF) - all of which affect the modeling of the $t\bar{t}$ signal. The dominant systematic, the jet energy scale (JES) uncertainty, is sub-divided into several components (‘‘levels’’) in order to more accurately accommodate our best estimate of the relevant correlations [11]. Additional categories have been added in order to accommodate specific types of correlations. Each uncertainty category is discussed below.

Statistical: The statistical uncertainty associated with the M_{top} determination.

iJES: The statistical uncertainty on the JES arising from the in-situ $W \rightarrow qq'$ calibration alone. Residual JES uncertainties, which arise from effects not considered in the in-situ calibration, are included in the Method category below.

aJES: This is specific to $D\bar{O}$ Run II and is only included here in order to be consistent with reference [10].

bJES: The systematic uncertainty specific to the modeling of b-jets. This includes uncertainties arising from variations in the semi-leptonic branching fraction, b-fragmentation modeling, and differences in the color flow between b-quark jets and light-quark. This is usually labeled ‘‘B Jet’’ for CDF Run II analyses.

cJES: The systematic uncertainty on the JES arising from the modeling of the out-of-cone corrections. This is the quadrature sum of the L7 and L8 JES uncertainties for CDF Run II analyses.

dJES: The systematic uncertainty on the JES arising from the relative corrections. This is the L1 JES uncertainties for CDF Run II analyses.

rJES: The systematic uncertainty on the JES arising from the modeling of the calorimeter response, the underlying event, and the multiple interaction corrections. This is the quadrature sum of the L4, L5, and L6 JES uncertainties for CDF Run II analyses.

Lepton P_T : The systematic uncertainty arising from uncertainties in the scale of the lepton transverse momentum measurements. This is an important uncertainty in the analysis with minimal dependence on the jet energy scale. It was not considered a source of systematic uncertainty in the Run I measurements.

Signal: The systematic uncertainty arising from uncertainties in the modeling of the $t\bar{t}$ signal including variations in the ISR, FSR, and PDF descriptions used to generate the $t\bar{t}$ Monte Carlo samples that calibrate each method.

Generator: The systematic uncertainty associated with variations of the physics model used to calibrate the fit methods and correlated across all measurements. It includes variations observed when substituting Pythia [12–14] (Run I and Run II) or ISAJET [15] (Run I) for HERWIG [16, 17] when modeling the $t\bar{t}$ signal.

Uranium noise and multiple interactions (UN/MI): This is specific to DØ Run I and is only included here in order to be consistent with reference [10].

Background: The systematic uncertainty arising from uncertainties in modeling the dominant background sources, including q^2 variations. This is the quadrature sum of the “Background Shape” and “Background normalization” uncertainties for most CDF Run II analyses.

Method: The systematic uncertainty arising from any source specific to a particular fit method, including the variations in B-tagging efficiency and the finite Monte Carlo statistics available to calibrate each method. This is the quadrature sum of the “Method”, “B-tag”, and “MC Statistics” categories for most CDF Run II analyses.

Color Reconnections (CR): The systematic uncertainty arising from a variation of the phenomenological description of color reconnection between final state particles [18]. This is obtained taking the difference between PYTHIA 6.4 tune “Apro” and PYTHIA 6.4 tune “ACRpro” that only includes a change in the color reconnection model. Monte Carlo generators which explicitly include CR models for hadron collisions have recently become available. This was not possible in Run I and these measurements do not include this source of systematic uncertainty.

Multiple Hadron Interactions (MHI): The systematic uncertainty arising from a mismodeling of the distribution of number of collision per bunch crossing due to the change in the collider instantaneous luminosity during data-taking. It has been separated from other sources to account for the fact that it is uncorrelated with DØ’s measurements.

These categories represent the current preliminary understanding of the various uncertainty categories and their correlations. We expect these to evolve as we continue to probe each method’s sensitivity to the various systematic sources with ever improving precision. Small variations in the assignment of uncertainties to the uncertainty categories and in the correlations assumed negligibly affect the combination.

III. INPUT MEASUREMENTS

For this combination we use seven measurements: three published Run I [1-3] results, two published Run II results [4,5] and two preliminary Run II results [6,7]. They are summarized in Table I. The correlations between the various inputs are described in the next section. Based on studies described in reference [19] the statistical correlation between the $L_{xy} + P_T^{lep}$ and LJT inputs is set to zero in the combination. Variations of ± 0.05 , which cover the full range observed in these studies, negligibly affect the combination reported here.

There are other CDF measurements in the LJT, DIL, HAD, and minimal-JES-dependence channels. We have chosen to include in this combination the analysis methods which yielded the best *expected* sensitivity for each channel. The channels themselves are statistically independent of each other and are treated as such.

For the $L_{xy} + P_T^{lep}$ measurement, we include the systematic uncertainty associated with the potential mis-modeling of the background decay length distribution in the “Method” category, as it is a source of uncertainty unique to this method.

For the Run I measurements, we back propagate the systematic uncertainty specific to B-jets (bJES) as determined in Run II and then correct the Run I absolute corrections (rJES) to keep the total JES uncertainty constant. We perform a series of cross-check to probe the robustness of the result. We propagate using the estimation of the color reconnection systematic performed in Run II back to Run I measurements as well, finding no noticeable difference in the combination. Another cross-check is to back-propagate the Lepton P_T systematic back to Run I results; again, no noticeable difference was found in the combination. Variations of the above assumptions were explored and found to negligibly affect the final value.

IV. CORRELATIONS

The following correlations are used when making the combination:

- The uncertainties in the Statistical, Method, and iJES categories are taken to be uncorrelated among the measurements.

Input	Run II Preliminary		Run II Published		Run I Published		
	LJT	DIL	HAD	$L_{xy}+P_T^{lep}$	LJT	DIL	HAD
M_{top}	173.00	170.56	174.8	175.30	176.1	167.4	186.0
Statistical	0.65	2.19	1.70	6.20	5.1	10.3	10.0
iJES	0.58	0.0	1.64	0.0	-	-	-
aJES	-	-	-	-	-	-	-
bJES	0.26	0.35	0.21	0.0	0.6	0.5	0.6
cJES	0.27	2.01	0.49	0.0	2.7	2.6	3.0
dJES	0.01	0.64	0.08	0.0	0.7	0.6	0.3
rJES	0.41	1.98	0.21	0.10	3.4	2.8	4.0
Lepton P_T	0.14	0.31	-	1.10	-	-	-
Signal	0.21	0.36	0.23	1.60	2.6	2.8	1.8
Generator	0.37	0.57	0.31	0.60	0.1	0.6	0.8
UN/MI	-	-	-	-	-	-	-
Background	0.34	0.27	0.35	1.60	1.3	0.3	1.7
Method	0.10	0.05	0.67	1.40	0.0	0.7	0.6
Color Reconnections	0.37	0.61	0.41	0.40	-	-	-
Multiple Hadron Interactions	0.10	0.27	0.17	0.70	-	-	-
Statistical	0.65	2.19	1.70	6.20	5.1	10.3	10.0
Systematics	1.06	3.09	1.99	3.10	5.3	4.9	5.7
Total	1.24	3.79	2.61	6.90	7.3	11.4	11.5

TABLE I: The measurements used to determine the CDF combined top-quark mass. All numbers are in units of GeV/c^2 . The uncertainty categories and their correlations are defined in the text. Some of the systematic sources were not accounted for in Run I measurements, due to them being negligible with respect to the actual measurements.

- The uncertainties in the aJES, dJES, Lepton P_T and Multiple Hadron Interaction categories are taken to be 100% correlated among all Run I and all Run II measurements, but uncorrelated between Run I and Run II.
- The uncertainties in the Background category are taken to be 100% correlated among all measurements in the same channel.
- The uncertainties in the bJES, cJES, rJES, Signal, Generator and Color Reconnection categories are taken to be 100% correlated among all measurements.

Using the inputs from Table I and the correlations specified here, the resulting matrix of total correlation coefficients is given in Table II.

To help quantify how sensitive the combination is to these assumptions we vary some of these correlations and redo the combination. We change all correlations that were set to be 100%, to a 50% value and find a shift in the combined M_{top} of less than $50 \text{ MeV}/c^2$, and a change in precision of less than $10 \text{ MeV}/c^2$. We also vary the statistical correlation for the two very different LJT analyses from 0% to 50%, and find a shift and change in precision less than $10 \text{ MeV}/c^2$.

	Run II Preliminary		Run II Published		Run I Published		
	LJ	DIL	HAD	$L_{xy}+P_T^{lep}$	LJT	DIL	HAD
LJT	1						
DIL	0.43	1					
HAD	0.19	0.21	1				
$L_{xy}+P_T^{lep}$	0.19	0.12	0.06	1			
LJT	0.36	0.48	0.15	0.16	1		
DIL	0.20	0.28	0.10	0.08	0.29	1	
HAD	0.23	0.35	0.12	0.07	0.32	0.19	1

TABLE II: The resulting matrix of total correlation coefficients used to determine the CDF combined top quark mass.

V. RESULTS

Using the measurements of Table I and the correlations of Section IV the CDF combined top mass is

$$M_{top} = 173.13 \pm 0.67 \text{ (stat)} \pm 0.95 \text{ (syst)} \text{ GeV}/c^2 \quad (1)$$

$$= 173.13 \pm 1.16 \text{ GeV}/c^2 \quad (2)$$

with $\chi^2/dof = 3.6/6$, which corresponds to a chi-squared probability of about 73%, indicating good agreement among all the input measurements. The combined value is $0.5 \text{ GeV}/c^2$ higher than the previous CDF combination [9], due to an upward shift in the most precise determination in the LJT channel [6]. It is also $0.1 \text{ GeV}/c^2$ higher than [6] due to the second most precise determination in the HAD channel [5] measuring a somewhat higher value.

This CDF-only combination has a relative precision of $\Delta M_{top}/M_{top} = 0.67\%$. The combined value presented here improves the precision by 6% with respect to the single most precise measurement [6]. This CDF combination improves the precision on M_{top} by 23% with respect to the previous CDF best determination [9].

The combination presented provides a value practically identical to the previous Tevatron best determination of M_{top} [10], improving the precision by 7%.

The method of reference [8] decomposes the total uncertainty into the contributions from the various uncertainty categories; the results can be found in Table III.

Results	CDF combined values
M_{top}	173.13
iJES	0.58
aJES	-
bJES	0.24
cJES	0.17
dJES	0.03
rJES	0.23
Lepton P_T	0.12
Signal	0.12
Generator	0.36
UM	-
Background	0.28
Fit	0.13
CR	0.39
MHI	0.11
Syst.	0.95
Stat.	0.67
Total	1.16

TABLE III: Summary of the Tevatron combined world average M_{top} . The error categories are described in the text. The total systematic uncertainty and the total uncertainty are obtained by adding the relevant contributions in quadrature. All numbers are in units of GeV/c^2 .

The corresponding pull and weight for each of the inputs are listed in Table IV.

	Run II Preliminary		Run II Published		Run I Published		
	LJT	DIL	HAD	L_{xy}	LJT	DIL	HAD
Pull	-0.29	-0.71	+0.71	+0.32	+0.41	-0.51	+1.12
Weight (%)	+92.3	-2.0	+13.7	-0.01	-2.8	-0.4	-0.8

TABLE IV: The pull and weight for each of the inputs used to determine the CDF combined top quark mass.

The input measurements, the output combination and the χ^2/dof are summarized in Figure 3.

The weights of some of the measurements are negative. In general, this situation can occur if the correlation between two measurements is larger than the ratio of their total uncertainties. This is indeed the case here. In these instances the less precise measurement will usually acquire a negative weight. While a weight of zero means that a particular




FIG. 1: Relative weights of the input measurements in the combination. The relative weights have been obtained dividing each measurement weight by the sum over all measurements of the modulo of their weights.

input is effectively ignored in the combination, a negative weight means that it affects the resulting central value and helps reduce the total uncertainty. See Reference [21] for further discussion of negative weights. To visualize the weight each measurement carries in the combination, weights have been divided by the sum of the modulo of all the input measurements weights. The result is shown in Figure 1.

Although the chi-squared from the combination of all measurements indicates that there is good agreement among them, and no input has an anomalously large pull, it is still interesting to also fit for the top mass in the LJT, DIL, and HAD channels separately. We use the same methodology and include the systematic correlations among the measurements as described in Section IV. The results are shown in Table V.

Using the expression in reference [20] we calculate the following chi-squares $\chi^2(LJT - DIL) = 1.48/1$, $\chi^2(LJT - HAD) = 0.73/1$, and $\chi^2(DIL - HAD) = 2.57/1$. These correspond to chi-squared probabilities of 23%, 39%, and 11%, respectively, and indicate that all channels are reasonably consistent with each other.

	fit value (GeV/c ²)	correlations		
		M(LJT)	M(DIL)	M(HAD)
M(LJT)	172.75 ± 1.20	1		
M(DIL)	169.01 ± 3.32	0.38	1	
M(HAD)	174.99 ± 2.57	0.19	0.22	1

TABLE V: The results of a fit to determine the top-quark mass in the three final states separately.

Using only the Run II measurements of Table I and the correlations of Section IV the CDF combined Run II top




FIG. 2: A summary of the input measurements utilized in this combination, and the resulting CDF combined top-quark mass. The preliminary results are the ones that carry the asterisk sign. All other results have been published.

mass is

$$M_{top} = 173.37 \pm 1.17 \text{ GeV}/c^2 \quad (3)$$

with $\chi^2/dof = 1.19/3$, which corresponds to a chi-squared probability of about 75%, indicating good agreement among all the input measurements. The combined value is $240 \text{ MeV}/c^2$ higher than the Run I plus Run II CDF combination, and has only $10 \text{ MeV}/c^2$ larger uncertainty.

VI. CONCLUSION

We have combined CDF Run I and Run II top-quark mass measurements from all three final states, HAD, LJT, and DIL to get a CDF combined top-quark mass of $173.13 \pm 1.16 \text{ GeV}/c^2$. This CDF-only combination has thus a relative precision of $\Delta M_{top}/M_{top} = 0.67\%$. This corresponds to a relative improvement of 23% with respect to the previous CDF combination [9] and to a relative improvement of 7% with respect to the previous Tevatron combination [10]. Rounding off to two significant digits, the combination yields $M_{top} = 173.1 \pm 1.2(\text{stat} + \text{syst}) \text{ GeV}/c^2$.




FIG. 3: Extrapolation of the top mass measurement result to current and future integrated luminosity. We obtain the average luminosity for the combination of 5 fb^{-1} by weighting the luminosity analyzed in each measurement by the relative weight it carries in the combination. The dashed line shows the extrapolation assuming the statistical and systematic uncertainties scale down with luminosity. The solid line uses the more conservative assumption that only the statistical uncertainty will scale down with luminosity. The plot on the left extrapolates from the results obtained using 680 pb^{-1} of data, while the plot on the right extrapolates from the most recent CDF combination.

Acknowledgments

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the World Class University Program, the National Research Foundation of Korea; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

-
- [1] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **80**, (1998) 2779;
 - [2] CDF Collaboration, F. Abe *et al.*, Phys. Rev. **D63**, (2001) 032003;
 - [3] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **79**, (1997) 1992;
 - [4] T. Aaltonen *et al.* [The CDF Collaboration], Phys. Rev. D **81** (2010) 032002;
 - [5] T. Aaltonen *et al.* [The CDF Collaboration], Phys. Rev. D **81** (2010) 052011;
 - [6] T. Aaltonen *et al.* [The CDF Collaboration], CDF Conference Note 10198;
 - [7] T. Aaltonen *et al.* [The CDF Collaboration], CDF Conference Note 10033;
 - [8] L. Lyons, D. Gibaut, and P. Clifford, Nucl. Instrum. Meth. **A270** (1988) 110;
A. Valassi, Nucl. Instrum. Meth. **A500** (2003) 391;
 - [9] M. Datta, D. Glenzinski, F. Margaroli, G. Velev [The CDF collaboration], CDF Conference Note 9714;
 - [10] Tevatron Electroweak Working Group, [The CDF and D0 Collaborations], arXiv:0903.2503;
 - [11] A. Bhatti *et al.*, Nucl. Instrum. Meth. A **566** (2006) 375;
 - [12] H.-U. Bengtsson and T. Sjostrand, Comput. Phys. Commun. **46** (1987) 43;
 - [13] T. Sjostrand, Comput. Phys. Commun. **82** (1994) 74;
 - [14] T. Sjostrand *et al.*, Comput. Phys. Commun. **135** (2001) 238;
 - [15] F. E. Paige and S. D. Protopopescu, BNL Reports 38034 and 38774 (1986) unpublished;
 - [16] G. Marchesini *et al.*, Comput. Phys. Commun. **67** (1992) 465;
 - [17] G. Corcella *et al.*, JHEP **01** (2001) 010;
 - [18] P. Z. Skands, arXiv:0905.3418;
 - [19] T. Aaltonen *et al.* [The CDF Collaboration], CDF Conference Note 8222;
 - [20] For two measurements, x and y , we calculate their consistency using $\chi^2 = (x-y)^2/\sigma_{x-y}^2$, where $\sigma_{x-y}^2 = \sigma_x^2 + \sigma_y^2 - 2\rho_{xy}\sigma_x\sigma_y$, where ρ_{xy} is the (x, y) correlation coefficient;
 - [21] L. Lyons, D. Gibaut, and P. Clifford, Nucl. Instrum. Meth. **A270** (1988) 110.