





Higgs searches and prospects at CDF




Pavel Murat (FNAL) for CDF collaboration



P. Murat, HCP-2002, Karlsruhe




Overview




- **Standard Model and Higgs boson**
- **CDF Run I searches of SM Higgs**
- **Run II improvements**
- **Run II prospects**
- **Higgs bosons in MSSM**
- **CDF Run I searches of MSSM Higgs bosons**
- **Run II prospects for MSSM Higgs searches**
- **Summary**



The Standard Model (SM) and the Higgs Boson




- SM extremely successful over the last 3 decades
 - Consistent with all the data
 - Predicted charm and top
- Higgs field is a key component of SM:
 - generates masses of all the particles
 - keeps the theory renormalizable at EW scale
- All constituents of SM, but the Higgs boson, observed experimentally
- Lower limit on the Higgs mass comes from direct searches
 - $M_H > 114.4 \text{ GeV}$ (LEP'2002, 95% CL)
- Global fit of EW data suggests upper bound:
 - $M_H < 193 \text{ GeV}$ (LEP'2002, 95% CL)






SM Higgs production at hadron colliders




- Cross sections for $M_H = 120$
 - Gluon fusion ($gg \rightarrow H$) : 0.7 pb
 - Associative production :
 - $qq \rightarrow WH$: 0.16 pb
 - $qq \rightarrow ZH$: 0.10 pb
- Smallest cross section measured so far:
 - $\sigma(pp \rightarrow tt) \sim 5\text{pb}$






SM Higgs : decays




- **‘Light’ Higgs: $M < 130$ GeV**
 - Dominant decay: $H \rightarrow b\bar{b}$
- **‘Heavy’ Higgs ($130 < M < 180$ GeV)**
 - $H \rightarrow W^*W$





Triggering and reconstruction




- $\sigma(p\text{-}p\text{bar}) \sim 70 \text{ mb}$
- For $\sigma(pp \rightarrow H) \sim 1 \text{ pb}$ **signal/background $\sim 10^{-10}$**
- **Need efficient trigger with high background rejection**

$M_H < 130 \text{ GeV}$

- **$pp \rightarrow H \rightarrow bb$** swamped with the QCD
- **$pp \rightarrow V(l\ell, l\nu, \nu\nu, qq)H(bb)$**
 - trigger: high-Pt leptons and/or missing E_t from W/Z, multi-jet events
 - Reconstruction:
 - Identify B-jets from Higgs decay (B-tagging)
 - Reconstruct M_H



Triggering and reconstruction




130 GeV < M(H) < 180 GeV


- **pp->H(W)->WW*(W)**
 - Signature: ll(l) + MET
 - Trigger:
 - High-Pt leptons from W's
 - Dileptons
 - same-sign dileptons
 - Reconstruction:
 - Can't reconstruct M_H (2 ν in the final state)
 - Counting experiment



Run I Search in $pp \rightarrow HZ \rightarrow llbb$




- Signature: 2 OS leptons (Z) + 2 jets
- The cleanest channel, smallest $\sigma \cdot BR$
- Trigger : high Pt ($Pt > 20 \text{ GeV}$) lepton
- Observed: 5 events
- Expected: 4 ± 1 events
- Irreducible QCD backgrounds dominate
 - $pp \rightarrow Zbb$




Cut	N_{ee}	$N_{\mu\mu}$
2 leptons	7029	3758
$76 < M_{ll} < 106$	5892	2880
2 or 3 15 GeV jets	105	54
$\cancel{E}_T < 50 \text{ GeV}$	105	53
≥ 1 SVX tag	2	3



Search in $pp \rightarrow WH \rightarrow l\nu b\bar{b}$ channel




- Signature: lepton+MET+ (≥ 2 jets)
- Trigger: high Pt lepton (e,mu)
- Lepton Pt > 20 GeV, $|\eta| < 1$
- Jet Et > 15 GeV, $|\eta| < 2$.
- MET > 20 GeV
- Require 1 or 2 jets to be B-tagged
- **Single tag**
 - Observed: 36 (22 e + 14 mu)
 - Expected: 30 ± 5
- **Double tag**
 - Observed: 6 (2 e + 4 mu)
 - Expected: 3.8 ± 0.7






Search in pp -> VH -> qq bb channel




- Signature: ≥ 4 jets
- Trigger: $\text{sum}(E_t) > 125 \text{ GeV}$
- Jet $E_t > 15 \text{ GeV}$, $|\eta| < 2.1$
- 2 jets B-tagged
- $P_t(bb) > 50 \text{ GeV}$
- Observed: 589 events
- Expected: 594 ± 30 , irreducible QCD background dominates






Search in $pp \rightarrow ZH \rightarrow \nu\nu b\bar{b}$ channel




- Signature: ≥ 2 jets + missing E_t
- Trigger: $MET > 40$ GeV
- Jet $ET > 15$ GeV and $|\eta| < 2$.
- 1 or 2 B-tagged jets
- **Single tag analysis:**
 - Observed: 40 events
 - Expected: 43 ± 5
- **Double tag analysis:**
 - Observed: 4 events
 - Expected: 4.9 ± 0.6





Summary of CDF Run I SM Higgs searches




- Searched for associative (W,Z) H production in 4 different channels

channel	L, pb^{-1}	N(observed)	N(bgr)
$llb\bar{b}$	106 ± 4	5	4.0 ± 1.0
$\nu\nu bb$	87 ± 4	4	4.9 ± 0.6
$l\nu bb$	106 ± 4	6	3.8 ± 0.7
$qqbb$	87 ± 4	589	594 ± 30


Benchmark number ($M = 130$ GeV):

$$\sigma(pp \rightarrow VH) * BR(H \rightarrow bb) < 7.4 \text{ pb @ 95\% CL}$$






Run II improvements





- Integrated luminosity: 2pb^{-1} (**x20**)/ $10\text{-}15\text{pb}^{-1}$ (**x100-150**)
- E_{cm} : **~40%** in $\sigma(\text{gg}\rightarrow\text{H})$, **~20%** in $\sigma(\text{qq}\rightarrow\text{WH})$
- better trigger efficiency for $\text{VH}\rightarrow\text{qqbb}$ (B-jet trigger)
- B-tagging
 - Acceptance for 2 jets (**~x2**)
 - Resolution in impact parameter (**~x1.2**)
- Jet energy resolution (**~x1.3**)
- Combine results with DZero (**x2**)
- **Overall improvement in $S/\sqrt{B} \sim 30$ is achievable**



Jet energy resolution, scale




- New jet correction algorithm: adds tracks and shower max detector information
- Resolution improved by 30-40% -> 15-20% in S/\sqrt{B}
 - **Same as the test beam results for single pions!**
- $Z \rightarrow b \bar{b}$: B-jet energy calibration at 90 GeV






Heavy SM Higgs ($130 \text{ GeV} < M < 180 \text{ GeV}$)




- Production: $gg \rightarrow H$, $qq \rightarrow HW$
- Highest rate channels:
 - $pp \rightarrow H \rightarrow W^* W^* \rightarrow ll\nu\nu$
 - $pp \rightarrow HW \rightarrow WW^*W^* \rightarrow llqqX$
- Trigger: high Pt leptons (same sign dileptons for WWW)
- Major backgrounds: W^+W^- and $t t$
- Run II Higgs workshop:
 - signal can be observed
 - cross section low
- $M_c = \sqrt{(M^2(l) + Pt^2(l))} + MET$




$M(H)[\text{GeV}]$	140	160	180
Signal [fb]	2.8	1.5	0.96
Backgrounds [fb]	44.	4.4	3.8
S/\sqrt{B} for 30 fb^{-1}	2.1	3.9	2.7



SM Higgs: projections for Run II




- Run IIa:
 - **With 2 fb^{-1} exclude SM Higgs for masses up to 120 GeV**
- Run IIb:
 - **with $10\text{-}15 \text{ fb}^{-1}$ exclude mass range up to $\sim 180 \text{ GeV}$**
- Observation (5σ) requires significantly larger statistics
- To reach maximal sensitivity CDF and D0 will need to combine their results in multiple search channels







Higgs bosons in MSSM




- 2 Higgs doublets
- 3 neutral (h,H,A), 2 charged (H^\pm) scalars
- 2 parameters: $\tan\beta = v_u/v_d$ and M_A
- Mass constraints
 - $M(h) < 135 \text{ GeV}$, $M(H^\pm) > M(W)$
- **Couplings to fermions may be enhanced:**
 - **Abb: $\gamma_5 \tan\beta$**
- large $\tan\beta$ within the reach of the Tevatron Run I ($\sim 100 \text{ pb}^{-1}$)



- LEP'2001 95% CL limits:
 - $M(h) > 91 \text{ GeV}$,
 - $M(A) > 91.9 \text{ GeV}$
 - $M(H^\pm) > 78.6 \text{ GeV}$




Higgs boson signatures in MSSM




- SM-like signatures
 - $qq \rightarrow VH \rightarrow Vbb$
 - Interpret results of SM Higgs searches in terms of MSSM parameters
- Signatures, enhanced in MSSM (large $\tan\beta$)
 - **$gg, qq \rightarrow \phi^0 bb \rightarrow bbbb$**
- New signatures
 - **$H^+ \rightarrow \tau\nu$**
- Large cross sections into the final states with tau leptons
 - Decays **$\phi^0 \rightarrow \tau\tau$ and $H^+ \rightarrow \tau\nu$**
- **Taus can be triggered on!**



CDF Run I Search for the neutral MSSM Higgs boson




- Signature: $pp \rightarrow Hbb \rightarrow bbbb$
- 4 jets $E_t > 15 \text{ GeV}$, $|\eta| < 2.1$
- 2 jets $E_t > 45\text{-}70 \text{ GeV}$
- 3 B-tags
- Reconstruct $M(bb)$
- $\epsilon \sim 0.4\text{-}0.5\%$ (trigger, b-tagging)
- Backgrounds: ($\sim 90\%$ irreducible) dominated by QCD






Run I Search for H^+ in top decays dilepton channel




$pp \rightarrow tt \rightarrow H^+(\tau\nu)bW(l\nu)b$

- Signature: $l(e,\mu) + \tau + \text{MET} + jj$
- e,μ : $P_t > 20 \text{ GeV}$, $|\eta| < 1$
- T : $P_t > 15 \text{ GeV}$, $|\eta| < 1$
- 2 jets $E_t > 10$, $|\eta| < 2$
- MET significance > 3
- Acceptance*BR $\sim 1\%$
- **Observe 4 events, expect 3.1 ± 0.5**
- Backgrounds dominated by SM $t\bar{t}b\bar{b}$ production






Run I search for charged Higgs in top decays hadronic channels




$p \bar{p} \rightarrow t \bar{t} \rightarrow H^+(\tau\nu)b(W/H^+)b$

- Include **$W \rightarrow qq$** to increase trigger acceptance, search for 2 signatures
 - **$\tau \text{ MET } b j$ and $\tau\tau \text{ MET}$**
- Observed: 7 events (7+0)
- Expected: 7.4 +/- 2 events
- Background dominated by fakes, W +jets (2nd largest) is ~20%






MSSM prospects: detection of τ leptons




- MSSM at $\tan\beta \gg 1$: detecting tau leptons is a priority
 - Trigger on hadronic tau decays
- Implemented single tau and ditau triggers (more details in the talk by John Smith)
 - Tau - missing Et (W, H+)
 - Tau - lepton (e, mu) (Z, h/H/A)
 - Tau - tau (Z, h/H/A)
- **W- $\rightarrow\tau\nu$ signal in Run II data taken with tau-MET trigger**





Summary



- Run I established tools and techniques for Higgs searches at hadron colliders
- CDF potential for detecting the Higgs has greatly increased in Run II - new detectors, new triggers, new calibration and analysis tools
- (re)learning how to use all of those
- After the 1st year of Run II don't have enough sensitivity to confront the Standard Model
- By the end of the year enough statistics to start exploring MSSM limits
- **Expect first results on Higgs searches in 2003**